

Generation of Bianchi type V cosmological models with varying Λ -term

ANIRUDH PRADHAN, *)

Department of Mathematics, Hindu Post-graduate College, Zamania-232 331, Ghazipur, India

ANIL KUMAR YADAV, LALLAN YADAV

Department of Physics, K.N.Govt. Post-graduate College, Gyanpur, Sant Ravidas Nagar, Bhadohi - 221 304, India

Received XXX

Bianchi type V perfect fluid cosmological models are investigated with cosmological term Λ varying with time. Using a generation technique (Camci *et al.*, 2001), it is shown that the Einstein's field equations are solvable for any arbitrary cosmic scale function. Solutions for particular forms of cosmic scale functions are also obtained. The cosmological constant is found to be decreasing function of time, which is supported by results from recent type Ia supernovae observations. Some physical aspects of the models are also discussed.

PACS: 98.80.Es, 98.80.-k

Key words: cosmology, variable cosmological constant, Bianchi type V universe

1 Introduction

The study of Bianchi type V cosmological models create more interest as these models contain isotropic special cases and permit arbitrary small anisotropy levels at some instant of cosmic time. This property makes them suitable as model of our universe. The homogeneous and isotropic Friedman-Robertson-Walker (FRW) cosmological models, which are used to describe standard cosmological models, are particular case of Bianchi type I, V and IX universes, according to whether the constant curvature of the physical three-space, $t = \text{constant}$, is zero, negative or positive. These models will be interesting to construct cosmological models of the types which are of class one. Present cosmology is based on the FRW model which is completely homogeneous and isotropic. This is in agreement with observational data about the large scale structure of the universe. However, although homogeneous but anisotropic models are more restricted than the inhomogeneous models, they explain a number of observed phenomena quite satisfactorily. This stimulates the research for obtaining exact anisotropic solution for Einstein's field equations (EFEs) as a cosmologically accepted physical models for the universe (at least in the early stages). Roy and Prasad [1] have investigated Bianchi type V universes which are locally rotationally symmetric and are of embedding class one filled with perfect fluid with heat conduction and radiation. Bianchi type V cosmological mod-

*) E-mail:pradhan@iucaa.ernet.in, acpradhan@yahoo.com

els have been studied by other researchers (Farnsworth [2], Maartens and Nel [3], Wainwright *et al.* [4], Collins [5], Meena and Bali [6], Pradhan *et al.* [7, 8]) in different context.

Models with a relic cosmological constant Λ have received considerable attention recently among researchers for various reasons (see Refs. [9 - 13] and references therein). Some of the recent discussions on the cosmological constant “problem” and on cosmology with a time-varying cosmological constant by Ratra and Peebles [14], Dolgov [15 - 17] and Sahni and Starobinsky [18] point out that in the absence of any interaction with matter or radiation, the cosmological constant remains a “constant”, however, in the presence of interactions with matter or radiation, a solution of Einstein equations and the assumed equation of covariant conservation of stress-energy with a time-varying Λ can be found. For these solutions, conservation of energy requires decrease in the energy density of the vacuum component to be compensated by a corresponding increase in the energy density of matter or radiation. Earlier researchers on this topic, are contained in Zeldovich [19], Weinberg [10] and Carroll, Press and Turner [20]. Recent cosmological observations by High- z Supernova Team and Supernova Cosmological Project (Garnavich *et al.* [21], Perlmutter *et al.* [22], Riess *et al.* [23], Schmidt *et al.* [24]) strongly favour a significant and positive Λ with the magnitude $\Lambda(G\hbar/c^3) \approx 10^{-123}$. Their findings arise from the study of more than 50 type Ia supernovae with redshifts in the range $0.10 \leq z \leq 0.83$ and suggest Friedman models with negative pressure matter such as a cosmological constant, domain walls or cosmic strings (Vilenkin [25], Garnavich *et al.* [21]). The main conclusion of these observations on magnitude and red-shift of type Ia supernova suggest that the expansion of the universe may be an accelerating one with a large function of cosmological density in the form of the cosmological Λ -term.

Several ansatz have been proposed in which the Λ term decays with time (see Refs. Gasperini [26], Freese *et al.* [27], Özer and Taha [13], Peebles and Ratra [28], Chen and Hu [29], Abdussattar and Viswakarma [30], Gariel and Le Denmat [31], Pradhan *et al.* [32]). Of the special interest is the ansatz $\Lambda \propto S^{-2}$ (where S is the scale factor of the Robertson-Walker metric) by Chen and Wu [29], which has been considered/modified by several authors (Abdel-Rahaman [33], Carvalho *et al.* [13], Waga [34], Silveira and Waga [35], Vishwakarma [36]).

In recent years, several authors (Hajj-Boutros [37], Hajj-Boutros and Sfeila [38], Ram [39], Mazumder [40] and Pradhan and Kumar [41]) have investigated the solutions of EFEs for homogeneous but anisotropic models by using some different generation techniques. Bianchi spaces $I - IX$ are useful tools in constructing models of spatially homogeneous cosmologies (Ellis and MacCallum [42], Ryan and Shepley [43]). From these models, homogeneous Bianchi type V universes are the natural generalization of the open FRW model which eventually isotropize. Recently Camci *et al.* [44] derived a new technique for generating exact solutions of EFEs with perfect fluid for Bianchi type V spacetime. In this paper, in what follows, we will discuss Bianchi type V cosmological models obtained by augmenting the energy-momentum tensor of a perfect fluid by a term that represents a variable cosmological constant varying with time, and later generalize the solutions of Refs. [39, 44]. This

paper is organized as follows: The field equations and the generation technique are presented in Section 2. We relate three of the metric variables by solving the off-diagonal component of EFEs, and find a second integral which is used to relate the remaining two metric variables. In Section 3, for the particular form of each metric variables, some solutions are presented separately and solutions of Camci *et al.* [44] and Ram [39] are shown to be particular cases of our solutions. Kinematical and dynamical properties of all solutions are also studied in this section. In Section 4, we give the concluding remarks.

2 Field equations and generation technique

In this section, we review the solutions obtained by Camci *et al.* [44]. The usual energy-momentum tensor is modified by addition of a term

$$T_{ij}^{(vac)} = -\Lambda(t)g_{ij}, \quad (1)$$

where $\Lambda(t)$ is the cosmological term and g_{ij} is the metric tensor. Thus the new energy-momentum tensor is

$$T_{ij} = (p + \rho)u_i u_j - pg_{ij} - \Lambda(t)g_{ij}, \quad (2)$$

where ρ and p , respectively, the energy and pressure of cosmic fluid, and u_i is the fluid four-velocity such that $u^i u_i = 1$.

We consider the spacetime metric of the spatially homogeneous Bianchi type V of the form

$$ds^2 = dt^2 - A^2(t)dx^2 - e^{2\alpha x} [B^2(t)dy^2 + C^2(t)dz^2], \quad (3)$$

where α is a constant. For the energy momentum tensor (2) and Bianchi type V spacetime (3), Einstein's field equations

$$R_{ij} - \frac{1}{2}Rg_{ij} = -8\pi T_{ij} \quad (4)$$

yield the following five independent equations

$$\frac{A_{44}}{A} + \frac{B_{44}}{B} + \frac{A_4 B_4}{AB} - \frac{\alpha^2}{A^2} = -8\pi(p + \Lambda), \quad (5)$$

$$\frac{A_{44}}{A} + \frac{C_{44}}{C} + \frac{A_4 C_4}{AC} - \frac{\alpha^2}{A^2} = -8\pi(p + \Lambda), \quad (6)$$

$$\frac{B_{44}}{B} + \frac{C_{44}}{C} + \frac{B_4 C_4}{BC} - \frac{\alpha^2}{A^2} = -8\pi(p + \Lambda), \quad (7)$$

$$\frac{A_4 B_4}{AB} + \frac{A_4 C_4}{AC} + \frac{B_4 C_4}{BC} - \frac{3\alpha^2}{A^2} = 8\pi(\rho - \Lambda), \quad (8)$$

$$\frac{2A_4}{A} - \frac{B_4}{B} - \frac{C_4}{C} = 0. \quad (9)$$

Here and in what follows the suffix 4 by the symbols A , B , C and ρ denote differentiation with respect to t . The Bianchi identity ($T^i_{;j} = 0$) takes the form

$$\rho_4 + (\rho + p)\theta = 0. \quad (10)$$

For complete determinacy of the system, we consider a perfect-gas equation of state

$$p = \gamma\rho, \quad 0 \leq \gamma \leq 1. \quad (11)$$

It is worth noting here that our approach suffers from a lack of Lagrangian approach. There is no known way to present a consistent Lagrangian model satisfying the necessary conditions discussed in this paper.

The physical quantities expansion scalar θ and shear scalar σ^2 have the following expressions:

$$\theta = u^i_{;i} = \frac{A_4}{A} + \frac{B_4}{B} + \frac{C_4}{C} \quad (12)$$

$$\sigma^2 = \frac{1}{2}\sigma_{ij}\sigma^{ij} = \frac{1}{3}\left[\theta^2 - \frac{A_4B_4}{AB} - \frac{A_4C_4}{AC} - \frac{B_4C_4}{BC}\right]. \quad (13)$$

Integrating Eq. (9) and absorbing the integration constant into B or C , we obtain

$$A^2 = BC \quad (14)$$

without any loss of generality. Thus, elimination of p from Eqs. (5) - (7) gives the condition of isotropy of pressures

$$2\frac{B_{44}}{B} + \left(\frac{B_4}{B}\right)^2 = 2\frac{C_{44}}{C} + \left(\frac{C_4}{C}\right)^2, \quad (15)$$

which on integration yields

$$\frac{B_4}{B} - \frac{C_4}{C} = \frac{k}{(BC)^{3/2}}, \quad (16)$$

where k is a constant of integration. Hence for the metric function B or C from the above first order differential Eq. (16), some scale transformations permit us to obtain new metric function B or C .

Firstly, under the scale transformation $dt = B^{1/2}d\tau$, Eq. (16) takes the form

$$CB_\tau - BC_\tau = kC^{-1/2}, \quad (17)$$

where subscript represents derivative with respect to τ . Considering Eq. (17) as a linear differential equation for B , where C is an arbitrary function, we obtain

$$(i)B = k_1C + kC \int \frac{d\tau}{C^{5/2}}, \quad (18)$$

where k_1 is an integrating constant. Similarly, using the transformations $dt = B^{3/2}d\tilde{\tau}$, $dt = C^{1/2}dT$, and $dt = C^{3/2}d\tilde{T}$ in Eq. (16), after some algebra we obtain respectively

$$(ii) B(\tilde{\tau}; k_2, k) = k_2 C \exp\left(k \int \frac{d\tilde{\tau}}{C^{3/2}}\right), \quad (19)$$

$$(iii) C(T; k_3, k) = k_3 B - kB \int \frac{dT}{B^{5/2}}, \quad (20)$$

and

$$(iv) C(\tilde{T}; k_4, k) = k_4 B \exp\left(k \int \frac{d\tilde{T}}{B^{3/2}}\right), \quad (21)$$

where k_2 , k_3 and k_4 are constants of integration. Thus choosing any given function B or C in cases (i), (ii), (iii) and (iv), one can obtain B or C and hence A from (14).

3 Generation of new solutions

We consider the following four cases:

3.1 Case (i): Let $C = \tau^n$, (n is a real number satisfying $n \neq \frac{2}{5}$).

In this case, Eq. (18) gives

$$B = k_1 \tau^n + \frac{2k}{2-5n} \tau^{1-3n/2} \quad (22)$$

and then from (14), we obtain

$$A^2 = k_1 \tau^{2n} + \frac{2k}{2-5n} \tau^{1-n/2}. \quad (23)$$

Hence the metric (3) reduces to the new form

$$ds^2 = (k_1 \tau^n + 2\ell \tau^{\ell_1}) [d\tau^2 - \tau^n dx^2] - e^{2\alpha x} \left[(k_1 \tau^n + 2\ell \tau^{\ell_1})^2 dy^2 + \tau^{2n} dz^2 \right], \quad (24)$$

where

$$\ell = \frac{k}{2-5n} \text{ and } \ell_1 = 1 - \frac{3n}{2}.$$

For this derived model (24), the pressure, energy density and cosmological constant are given by

$$\begin{aligned} 8\pi(p + \Lambda) = & (k_1 \tau^n + 2\ell \tau^{\ell_1})^{-3} \left[-2k_1^2 n(n-1) \tau^{2n-2} - k_1 \ell n(10-13n) \tau^{-(\ell_1+2n)} \right. \\ & \left. - \frac{\ell^2(4+4n-11n^2)}{2} \tau^{-3n} \right] + \alpha^2 \tau^{-n} (k_1 \tau^n + 2\ell \tau^{\ell_1})^{-1}, \end{aligned} \quad (25)$$

$$8\pi(\rho - \Lambda) = (k_1\tau^n + 2\ell\tau^{\ell_1})^{-3} \left[3k_1^2 n^2 \tau^{2n-2} + 3k_1 \ell n(2-n)\tau^{-(\ell_1+2n)} + \frac{\ell^2(4+4n-11n^2)}{2} \tau^{-3n} \right] - 3\alpha^2 \tau^{-n} (k_1\tau^n + 2\ell\tau^{\ell_1})^{-1}. \quad (26)$$

Eq. (25), with the use of (11) and (26), reduces to

$$8\pi(1+\gamma)\rho = (k_1\tau^n + 2\ell\tau^{\ell_1})^{-3} \left[k_1^2 n(n+1)\tau^{2n-2} + 2k_1 \ell n(5n-2)\tau^{-(\ell_1+2n)} \right] - 2\alpha^2 \tau^{-n} (k_1\tau^n + 2\ell\tau^{\ell_1})^{-1} \quad (27)$$

Eliminating $\rho(t)$ between (26) and (27), we obtain

$$8\pi(1+\gamma)\Lambda = (k_1\tau^n + 2\ell\tau^{\ell_1})^{-3} \left[-k_1^2 n\{(2+3\gamma)n-2\}\tau^{2n-2} - k\ell n\{10-13n+3\gamma(2-n)\}\tau^{-(\ell_1+2n)} - \frac{(1+\gamma)\ell^2(4+4n-11n^2)}{2} \tau^{-3n} \right] + (1+3\gamma)\alpha^2 \tau^{-n} (k_1\tau^n + 2\ell\tau^{\ell_1})^{-1}. \quad (28)$$

From Eq. (28), we observe that at the time of early universe the cosmological constant (Λ) increases rapidly during a very short time and then it decreases as time increases (see Figure 1). We also observe that the value of Λ is small and positive at late times which is supported by recent type Ia supernovae observations [22 - 24].

The metric (24) is a four-parameter family of solutions to EFEs with a perfect fluid. Using the scale transformation $dt = B^{1/2}d\tau$ in Eqs. (12) and (13) for this case, the scalar expansion θ and the shear σ have the expressions:

$$\theta = 3 \left[k_1 n \tau^{n-1} + \frac{\ell(2-n)}{2} \tau^{-3n/2} \right] (k_1\tau^n + 2\ell\tau^{\ell_1})^{-3/2} \quad (29)$$

$$\sigma = \frac{1}{2} k \tau^{-3n/2} (k_1\tau^n + 2\ell\tau^{\ell_1})^{-3/2} \quad (30)$$

Eqs. (29) and (30) lead to

$$\frac{\sigma}{\theta} = \frac{k}{6} \left[k_1 n \tau^{n-\ell_1} + \frac{\ell(2-n)}{2} \right]^{-1} \quad (31)$$

Now, we consider four subcases for the parameters Λ , n , k , k_1 whether zero or not. In subcase $\Lambda = 0$, metric (24) with expressions p , ρ , θ and σ for this model are same as that of solution (18) of Camci *et al.* [44].

In subcase $\Lambda = 0$, $n = 0$, after a suitable inverse time transformation, we find that

$$ds^2 = dt^2 - K(t+t_0)^{2/3} dx^2 - e^{2\alpha x} \left[K_1(t+t_0)^{4/3} dy^2 + dz^2 \right], \quad (32)$$




Fig. 1. The plot of cosmological constant Λ versus time τ for the model (24) in Section 3.1 Case(i) with parameters $\alpha = 1$, $n = 0.25$, $k = 1$, $k_1 = 1$, $\gamma = 0.5$.

where t_0 is a constant of integration and $K_1 = (3k/2)^{2/3}$. The expressions p , ρ , θ and σ for this model are not given here, since it is observed that the physical properties of this one are same as that of the solution (24) of Ram [39].

In subcase $\Lambda = 0$, $k = 0$, after inverse time transformation and rescaling, the metric (24) reduces to

$$ds^2 = dt^2 - K_2(t + t_1)^{\frac{4n}{n+2}} [dx^2 + e^{2\alpha x}(dy^2 + dz^2)], \quad (33)$$

where t_1 is a constant of integration and $K_2 = \left(\frac{n+2}{2}\right)^{\frac{4n}{n+2}}$. For this solution, when $n = 1$ and $\alpha = 0$, we obtain Einstein and de Sitter [45] dust filled universe. For $K_2 = 1$, $t_1 = 0$ and $n = \frac{2m}{(2-m)}$, where m is a parameter in Ram's paper [39], the solution (33) reduces to the metric (14) of Ram [39]. In later case, if also $\alpha = 0$, then we get the Minkowski spacetime.

Now, in subcase $\Lambda = 0$, $k_1 = 0$, after some algebra the metric (24) takes the form

$$ds^2 = dt^2 - 2\ell K_3(t + t_2)^{2/3} [dx^2 + e^{2\alpha x}(at^{m_1}dy^2 + a^{-1}t^{-m_1}dz^2)], \quad (34)$$

where t_2 is a constant, $at^{m_1} = 2\ell K_3^{\frac{2-5n}{2-n}}(t+t_2)^{\frac{2(2-5n)}{3(2-n)}}$ and $K_3 = \left[\frac{3(2-n)}{4\sqrt{2\ell}}\right]^{2/3}$. For $t_2 = 0$, $k = \frac{2}{3}$ and $n = 0$ from (34), we obtain that the solution (24) of Ram [39].

Some Physical aspects of model :

The model (24) has barrel singularity at $\tau = \tau_0$ given by

$$\tau_0 = \left[\frac{k_1(5n-2)}{2k} \right]^{\frac{2}{(2-5n)}},$$

which corresponds to $t = 0$. For $n \neq 2/5$ from (24), it is observed that at the singularity state $\tau = \tau_0$, p , ρ , Λ , θ and σ are infinitely large. At $t \rightarrow \infty$, which corresponds to $\tau \rightarrow \infty$ for $n < 2/5$ and $k > 0$, or $\tau \rightarrow 0$ for $n > 2/5$ and $k < 0$, p , ρ , Λ , θ and σ vanish. Therefore, for $n \neq 2/5$, the solution (24) represents an anisotropic universe exploding from $\tau = \tau_0$, i.e. $t = 0$, which expands for $0 < t < \infty$. We also find that the ratio σ/θ tends to a finite limit as $t \rightarrow \infty$, which means that the shear scalar does not tend to zero faster than the expansion. Hence the model does not approach isotropy for large values of t .

In subcase $\Lambda = 0$, $k = 0$, the ratio (31) tends to zero, then the model approaches isotropy i.e. shear scalar σ goes to zero. For the model (33), p and ρ tends to zero as $t \rightarrow \infty$; the model would give an essentially empty universe at large time. The dominant energy condition given by Hawking and Ellis [46] requires that

$$\rho + p \geq 0, \quad \rho + 3p \geq 0 \quad (35)$$

Thus, we find for the model (33) that $n(2-n) \geq 0$. Hence for the values $0 \leq n \leq 2$, the universe (33) satisfies the strong energy condition i.e. $\rho + 3p \geq 0$. Also this model is shear-free and expanding.

In subcase $\Lambda = 0$, $k_1 = 0$, for $n \neq 2/5, 2$, it is observed from relations (25) - (30) that p , ρ , θ and σ are infinitely large at the singularity state $t = -t_2$. When $t \rightarrow \infty$, these quantities vanish. We also find that the ratio σ/θ is a constant. This shows that the cosmological model (34) does not approach isotropy for large value of t . In this model the dominant energy conditions (35) are then verified for $6 - 5n - 25n^2 \geq 0$. Since $n \neq 2/5$, the model (34) satisfies the strong energy condition for $-3/5 \leq n \leq 2/5$.

In each of subcases, all the obtained solutions (32), (33) and (34) satisfy the Bianchi identity given in Eq. (10).

3.2 Case (ii): Let $C = \tilde{\tau}^n$, (n is a real number satisfying $n \neq 2/3$).

In this case Eq. (19) gives

$$B = k_2 \tilde{\tau}^n \exp(M \tilde{\tau}^{\ell_1}) \quad (36)$$

and from (14), we obtain

$$A^2 = k_2 \tilde{\tau}^{2n} \exp(M \tilde{\tau}^{\ell_1}) \quad (37)$$

where $M = \frac{k}{\ell_1}$. Hence the metric (3) reduces to the form

$$ds^2 = \tilde{\tau}^{4(1-\ell_1)/3} \left[\tilde{\tau}^{2(1-\ell_1)/3} e^{3M\tilde{\tau}^{\ell_1}} d\tilde{\tau}^2 - e^{M\tilde{\tau}^{\ell_1}} dx^2 - e^{2\alpha x} \left(e^{2M\tilde{\tau}^{\ell_1}} dy^2 + dz^2 \right) \right], \quad (38)$$

where the constant k_2 is taken, without any loss of generality, equal to 1. This metric is a three-parameter family of solutions to EFEs with a perfect fluid. For the above model, the distribution of matter and nonzero kinematical parameters are obtained as

$$8\pi(p + \Lambda) = 2n\tilde{\tau}^{2(\ell_1-2)} + 3nk\tilde{\tau}^{3\ell_1-4} + \frac{k^2}{2}\tilde{\tau}^{4(\ell_1-1)} + \alpha^2\tilde{\tau}^{4(\ell_1-1)/3}e^{-3M\tilde{\tau}^{\ell_1}}, \quad (39)$$

$$8\pi(\rho - \Lambda) = 3n^2\tilde{\tau}^{2(\ell_1-2)} + 3nk\tilde{\tau}^{3\ell_1-4} + \frac{k^2}{2}\tilde{\tau}^{4(\ell_1-1)} - 3\alpha^2\tilde{\tau}^{4(\ell_1-1)/3}e^{-3M\tilde{\tau}^{\ell_1}}. \quad (40)$$

With the use of Eq. (11) in (39) and (40), we obtain

$$8\pi(1 + \gamma)\rho = n(2 + 3n)\tilde{\tau}^{2(\ell_1-2)} + 6nk\tilde{\tau}^{3\ell_1-4} + k^2\tilde{\tau}^{4(\ell_1-1)} - 2\alpha^2\tilde{\tau}^{4(\ell_1-1)/3}e^{-3M\tilde{\tau}^{\ell_1}}. \quad (41)$$

Eliminating $\rho(t)$ between (40) and (41), we get

$$8\pi(1 + \gamma)\Lambda = n(2 - 3n\gamma)\tilde{\tau}^{2(\ell_1-2)} + 3nk(1 - \gamma)\tilde{\tau}^{3\ell_1-4} + \frac{k^2}{2}(1 - \gamma)\tilde{\tau}^{4(\ell_1-1)} + \alpha^2(1 - 3\gamma)\tilde{\tau}^{4(\ell_1-1)/3}e^{-3M\tilde{\tau}^{\ell_1}}. \quad (42)$$

From Eq. (42), we observe that the cosmological constant (Λ) in the model (38) is a decreasing function of time (see Figure 2). We also observe that the value of Λ approaches a small and positive value at late times which is supported by recent type Ia supernovae observations [21 - 24].

The scalar of expansion θ and the shear σ are obtained as

$$\theta = 3 \left[n\tilde{\tau}^{\ell_1-2} + \frac{k}{2}\tilde{\tau}^{2(\ell_1-1)} \right], \quad (43)$$

$$\sigma = \frac{k}{2}\tilde{\tau}^{2(\ell_1-1)}e^{-3M\tilde{\tau}^{\ell_1}}, \quad (44)$$

From Eqs. (43) and (44), we have

$$\frac{\sigma}{\theta} = \frac{k}{6 \left(n\tilde{\tau}^{-\ell_1} + \frac{k}{2} \right)}. \quad (45)$$




Fig. 2. The plot of cosmological constant Λ versus time $\tilde{\tau}$ for the model (38) in Section 3.2 Case(ii) with parameters $\alpha = 1$, $n = 0.25$, $k = 1$, $\gamma = 0.5$.

In subcase $\Lambda = 0$, metric (38) with expressions p , ρ , θ and σ for this model are same as that of solution (27) of Camci *et al.* [44].

In sub-case $\Lambda = 0$, $\ell_1 = 1$ (i.e. $n = 0$), we find a similar solution to (32), and hence this subclass is omitted. For $k = 0$, the ratio (45) is zero and hence there is no anisotropy.

After a suitable coordinate transformation, the metric (38) can be written as

$$ds^2 = dt^2 - K_4(t + t_3)^{2\ell_1} [dx^2 + e^{2\alpha x}(dy^2 + dz^2)], \quad (46)$$

where t_3 is a constant and $K_4 = [2/(2 - 3M_1)]^{2M_1}$, $M_1 = \frac{2n}{2+3n} \neq 2/3$, where M_1 is a new parameter. When $M_1 = 0$ and $\ell_1 = 0$, from (46), we get the solution (12) of Ram [39].

Some physical aspects of model:

The model has singularity at $\tilde{\tau} \rightarrow -\infty$ for $\ell_1 > 0$ or $\tilde{\tau} \rightarrow 0$ for $\ell_1 < 0$, which

corresponds to $t \rightarrow 0$. It is a point type singularity for $\ell_1 > 0$ whereas it is a cigar or a barrel singularity according as $\ell_1 < 0$. At $t \rightarrow \infty$, which correspond to $\tilde{\tau} \rightarrow \infty$ for $\ell_1 > 0$ or $\tilde{\tau} \rightarrow 0$ for $\ell_1 < 0$, from Eqs. (39) - (45), we obtain that for $\ell_1 > 0$, $p, \rho \rightarrow 0$, and $\sigma, \theta \rightarrow 0$ ($k > 0$), $-\infty$ ($k < 0$); for $\ell_1 < 0$, similar the above ones. Then, clearly, for a realistic universe, it must be fulfill as $\tilde{\tau} \rightarrow -\infty$, n and k are positive and ℓ_1 is an odd positive number; as $\tilde{\tau} \rightarrow 0$, k is positive, and ℓ_1 an even negative number. Also, since $\lim_{t \rightarrow \infty} \frac{\sigma}{\theta} \neq 0$ therefore, this model does not approach isotropy for large values of t .

In subcase $k = 0$ for the metric (46), the pressure, density and cosmological constant are given by

$$8\pi(p + \Lambda) = \frac{\ell_1(2 - 3\ell_1)}{(t + t_3)^2} + \frac{\alpha^2}{K_4(t + t_3)^{2\ell_1}}, \quad (47)$$

$$8\pi(\rho - \Lambda) = \frac{3\ell_1}{(t + t_3)^2} - \frac{3\alpha^2}{K_4(t + t_3)^{2\ell_1}}. \quad (48)$$

Eq. (47), with the use of (11) and (48), reduces to

$$8\pi(1 + \gamma)\rho = \frac{2\ell_1}{(t + t_3)^2} - \frac{2\alpha^2}{K_4(t + t_3)^{2\ell_1}}. \quad (49)$$

Eliminating $\rho(t)$ between (48) and (49), we obtain

$$8\pi(1 + \gamma)\Lambda = \frac{\ell_1[2 - 3\ell_1(1 + \gamma)]}{(t + t_3)^2} + \frac{\alpha^2(1 + 3\gamma)}{K_4(t + t_3)^{2\ell_1}}. \quad (50)$$

When $\Lambda = 0$, the pressure and energy density are same as that of given in Eq. (44) of paper Camci *et al.* [44]. In this case, the weak and strong energy conditions (35) for this solution are identically satisfied when $\ell_1(1 - \ell_1) \geq 0$ i.e. $0 \leq \ell_1 \leq 1$. This model is shear-free and expanding with $\theta = \frac{3\ell_1}{(t + t_3)}$.

3.3 Case (iii) : Let $B = T^n$ (n is a real number).

In this case Eq. (20) gives

$$C = k_3 T^n - 2\ell T^{\ell_1} \quad (51)$$

and then from (14), we obtain

$$A^2 = k_3 T^{2n} - 2\ell T^{\ell_1 + n} \quad (52)$$

Hence the metric (3) takes the new form

$$ds^2 = (k_3 T^n - 2\ell T^{\ell_1}) [dt^2 - T^n dx^2] - e^{2\alpha x} [T^{2n} dy^2 + (k_3 T^n - 2\ell T^{\ell_1})^2 dz^2] \quad (53)$$

For four-parameters family of solution (53), the physical and kinematical quantities are given by

$$8\pi(p + \Lambda) = \left[-\frac{\ell^2}{2}(11n^2 - 4n - 4)T^{-3n} + \ell k_3 n(13n - 10)T^{\ell_1+n} - 2k_3^2 n(n-1)T^{2n-2} \right] (k_3 T^n - 2\ell T^{\ell_1})^{-3} + \alpha^2 T^{-n} (k_3 T^n - 2\ell T^{\ell_1})^{-1}, \quad (54)$$

$$8\pi(\rho - \Lambda) = \left[-\frac{\ell^2(11n^2 - 4n - 14)}{2}T^{-3n} - 3\ell k_3 n(2-n)T^{\ell_1+n} + 3k_3^2 n^2 T^{2n-2} \right] (k_3 T^n - 2\ell T^{\ell_1})^{-3} - 3\alpha^2 T^{-n} (k_3 T^n - 2\ell T^{\ell_1})^{-1}. \quad (55)$$

Eq. (54), with the use of (11) and (55), reduces to

$$8(1 + \gamma)\rho = \left[\ell^2(9 + 4n - 11n^2)T^{-3} + 16\ell k_3 n(n-1)T^{\ell_1+n} + k_3^2 n(2n+1)T^{2n-2} \right] (k_3 T^n - 2\ell T^{\ell_1})^{-3} - 2\alpha^2 T^{-n} (k_3 T^n - 2\ell T^{\ell_1})^{-1}. \quad (56)$$

Eliminating $\rho(t)$ between (55) and (56), we have

$$8(1 + \gamma)\Lambda = \left[-\frac{\ell^2}{2}\{n(11n-1)(1-\gamma) - 2(2-7\gamma)\}T^{-3} + \ell k_3 n\{13n-10+3(2-n)\gamma\}T^{\ell_1+n} - k_3^2 n\{2(n-1)-3n\gamma\}T^{2n-2} \right] (k_3 T^n - 2\ell T^{\ell_1})^{-3} + \alpha^2(1+3\gamma)T^{-n} (k_3 T^n - 2\ell T^{\ell_1})^{-1}. \quad (57)$$

From Eq. (57), we observe that the cosmological constant (Λ) in the model (53) decreases as time increases. We also observe that the value of Λ approaches a small and positive at late times which is supported by recent type Ia supernovae observations (21) – (24). Figure 3 clearly shows this behaviour of Λ as decreasing function of time.

The scale of expansion and the shear are obtained as

$$\theta = 3 \left[\frac{\ell(n-2)}{2}T^{-3n/2} + k_3 n T^{n-1} \right] (k_3 T^n - 2\ell T^{\ell_1})^{-3/2}, \quad (58)$$

$$\sigma = \frac{kT^{-3n/2}}{2} (k_3 T^n - 2\ell T^{\ell_1})^{-3/2}. \quad (59)$$

From (58) and (59), we get

$$\frac{\sigma}{\theta} = \frac{k}{6} \left[k_3 n T^{-\ell_1+n} + \frac{\ell(n-2)}{2} \right]^{-1}. \quad (60)$$




Fig. 3. The plot of cosmological constant Λ versus time T for the model (53) in Section 3.3 Case(iii) with parameters $\alpha = 1$, $n = 1$, $k = 1$, $k_3 = 1.5$, $\gamma = 0.5$.

In subcase $\Lambda = 0$, metric (53) with expressions p , ρ , θ and σ for this model are same as that of solution (34) of Camci *et al.* [44].

In subcase $\Lambda = 0$, $n = 0$, after an inverse transformation, metric (53) reduces to the form

$$ds^2 = dt^2 - K_5(t + t_4)^{2/3} dx^2 - e^{2\alpha x} \left[dy^2 + K_5^2(t + t_4)^{4/3} dz^2 \right], \quad (61)$$

where t_4 is an integrating constant. This model is different from the model (32) by a change of scale.

In subcase $\Lambda = 0$, $k = 0$, same model as (33) is obtained.

Further in subcase $\Lambda = 0$, $k_3 = 0$, we see that the metric (53) takes the form

$$ds^2 = dt^2 - 2\ell K_6(t + t_5)^{2/3} \left[dx^2 + e^{2\alpha x} (bt^{m_2} dy^2 + b^{-1}t^{-m_2} dz^2) \right], \quad (62)$$

where t_5 is a constant, $bt^{m_2} = 2\ell K_6^{\frac{2-5n}{2-n}}(t + t_5)^{\frac{2(2-5n)}{3(2-n)}}$ and $K_6 = \left[\frac{3(2-n)}{4\sqrt{2\ell}}\right]^{2/3}$. This metric is only different from (33) by a change of sign. Also, in each of subcase the physical and kinematical properties of obtained metric are same as that of case(i). Therefore, we do not consider here them.

3.4 Case (iv) : Let $B = \tilde{\tau}^n$, where n is any real number.

In this case Eq. (21) gives

$$C = k_4 \tilde{\tau}^n \exp\left(\frac{k}{\ell_1} \tilde{\tau}^{\ell_1}\right) \quad (63)$$

and then from (14), we obtain

$$A^2 = k_4 \tilde{\tau}^{2n} \exp\left(\frac{k}{\ell_1} \tilde{\tau}^{\ell_1}\right) \quad (64)$$

Hence the metric (3) reduces to

$$ds^2 = \tilde{\tau}^{2n} \exp\left(\frac{k}{\ell_1} \tilde{\tau}^{\ell_1}\right) \left[\tilde{\tau}^n \exp\left(\frac{2k}{\ell_1} \tilde{\tau}^{\ell_1}\right) - dx^2 \right] - e^{2\alpha x} \left[dy^2 + \exp\left(\frac{2k}{\ell_1} \tilde{\tau}^{\ell_1}\right) - dz^2 \right], \quad (65)$$

where, without any loss of generality, the constant k_4 is taken equal to 1. Expressions for physical and kinematical parameters for the model (65) are not given here, but it is observed that the properties of the metric (65) are same as that of the solution (38), i.e. the case (ii).

4 Concluding remarks

In this paper we have described a new exact solutions of EFES for Bianchi type V spacetime with a perfect fluid as the source of matter and cosmological term Λ varying with time. Using a generation technique followed by Camci *et al.* [44], it is shown that the Einstein's field equations are solvable for any arbitrary cosmic scale function. Starting from particular cosmic functions, new classes of spatially homogeneous and anisotropic cosmological models have been investigated for which the fluids are acceleration and rotation free but they do have expansion and shear. For $\alpha = 0$ in the metric (1), we obtained metrics as LRS Bianchi type I model (Hajj-Boutros [37], Hajj-Boutros and Sfeila [38], Ram [39], Mazumder [40] and Pradhan and Kumar [41]). It is also seen that the solutions obtained by Camci *et al.* [44] and Ram [39] are particular of our solutions.

The cosmological constants in all models given in Section 3 are decreasing functions of time and they all approach a small positive value at late times which are

supported by the results from recent supernova observations recently obtained by the High-z Supernova Team and Supernova Cosmological Project (Garnavich *et al.* [21], Perlmutter *et al.* [22], Riess *et al.* [23], Schmidt *et al.* [24]).

One of the authors (A. Pradhan) would like to thank the Inter-University Centre for Astronomy and Astrophysics, Pune, India for providing facility under Associateship Programme where part of this work was carried out. Authors are also thankful to C. S. Stalin for his help in plotting the figures.

References

- [1] S.R. Roy and A. Prasad: Gen. Rel. Grav. **26** (1994) 939.
- [2] D.L. Farnsworth: J. Math. Phys. **8** (1967) 2315.
- [3] R. Maartens and S.D. Nel: Comm. Math. Phys. **59** (1978) 273
- [4] J. Wainwright, W.C.W. Ince and B.J. Marshman: Gen. Rel. Grav. **10** (1979) 259.
C. G. Hewitt and J. Wainwright, Phys. Rev. D **46** (1992) 4242.
- [5] C.B. Collins: Comm. Math. Phys. **39** (1974) 131.
- [6] B.L. Meena and R. Bali: Pramana - journal of phys. **62** (2004) 1007).
- [7] A. Pradhan and A. Rai: Ast. Sp. Sc. **291** (2004) 149.
- [8] A. Pradhan, L. Yadav and A.K. Yadav: Czech. J. Phys. **54** (2004) 487.
- [9] S. Weinberg: Rev. Mod. Phys. **61** (1989) 1.
- [10] S. Weinberg: *Gravitation and Cosmology*, Wiley, New York, 1972.
- [11] J.A. Frieman and I. Waga: Phys. Rev. D **57** (1998) 4642.
- [12] R. Carlberg *et al.*: Astrophys. J. **462** (1996) 32.
- [13] M. Özer and M.O. Taha: Nucl. Phys. B **287** (1987) 776;
K. Freese, F.C. Adams, J.A. Frieman and E. Motta: *ibid.* B **287** (1987) 1797;
J.C. Carvalho, J.A.S. Lima and I. Waga: Phys. Rev. D **46** (1992) 2404;
V. Silveira and I. Waga: *ibid.* D **50** (1994) 4890.
- [14] B. Ratra and P.J.E. Peebles: Phys. Rev. D **37** (1988) 3406.
- [15] A.D. Dolgov: in *The Very Early Universe*, eds. G.W. Gibbons, S.W. Hawking and S.T.C. Siklos, Cambridge University Press, 1983.
- [16] A.D. Dolgov, M.V. Sazhin and Ya.B. Zeldovich: *Basics of Modern Cosmology*, Editions Frontiers, 1990.
- [17] A.D. Dolgov: Phys. Rev. D **55** (1997) 5881.
- [18] V. Sahni and A. Starobinsky: Int. J. Mod. Phys. D **9** (2000) 373.
- [19] Ya.B. Zeldovich: Sov. Phys.-Uspekhi **11** (1968) 381.
- [20] S.M. Carroll, W.H. Press and E.L. Turner: Ann. Rev. Astron. Astrophys. **30** (1992) 499.
- [21] P. M. Garnavich *et al.*: Astrophys. J. **493** (1998a) L53 (astro-ph/9710123); Astrophys. J. **509** (1998b) 74 (astro-ph/9806396).

- [22] S. Perlmutter *et al.*: *Astrophys. J.* **483** (1997) 565 (astro-ph/9608192); *Nature* **391** (1998) 51 (astro-ph/9712212); *Astrophys. J.* **517** (1999) 565.
- [23] A. G. Riess *et al.*: *Astron. J.* **116** (1998) 1009 (astro-ph/98052201).
- [24] B. P. Schmidt *et al.*: *Astrophys. J.* **507** (1998) 46 (astro-ph/98052200).
- [25] A. Vilenkin: *Phys. Rep.* **121** (1985) 265.
- [26] M. Gasperini: *Phys. Lett. B* **194** (1987) 347; *Class. Quant. Grav.* **5** (1988) 521.
- [27] K. Freese, F.C. Adams, J.A. Friemann and E. Mottolla: *Nucl. Phys. B* **287** (1987) 797.
- [28] P.J.E. Peebles and B. Ratra: *Astrophys. J.* **325** (1988) L17.
- [29] W. Chen and Y.S. Wu: *Phys. Rev. D* **41** (1990) 695.
- [30] Abdussattar and R.G. Vishwakarma: *Pramana - J. Phys.* **47** (1996) 41.
- [31] J. Gariel and G.Le Denmat: *Class. Quant. Grav.* **16** (1999) 149.
- [32] A. Pradhan and V.K. Yadav: *Int. J. Mod. Phys. D* **11** (2002) 893;
A. Pradhan and I. Aotemshi: *Int. J. Mod. Phys. D* **11** (2002) 1419.
- [33] A.-M. M. Abdel-Rahaman: *Gen. Rel. Grav.* **22** (1990) 655; *Phys. Rev. D* **45** (1992) 3492.
- [34] I. Waga: *Astrophys. J.* **414** (1993) 436.
- [35] V. Silveira and I. Waga: *Phys. Rev. D* **50** (1994) 4890.
- [36] R. G. Vishwakarma: *Class. Quant. Grav.* **17** (2000) 3833.
- [37] J. Hajj-Boutros: *Lecture Notes in Physics , Gravitation, Geometry and Relativistic Physics* Vol. 212, p. 51, Springer, Berlin (1984); *J. Math. Phys.* **26** (1985) 2297; *Class. Quant. Grav.* **3** (1986) 311.
- [38] J. Hajj-Boutros and J. Sfeila: *Int. J. Theor. Phys.* **26** (1987) 97.
- [39] S. Ram: *Int. J. Theor. Phys.* **29** (1990) 901; *Gen. Rel. Grav.* **21** (1989) 697.
- [40] A. Mazumder: *Gen. Rel. Grav.* **26** (1994) 307.
- [41] A. Pradhan and A. Kumar: *Int. J. Mod. Phys. D* **10** (2001) 291.
- [42] G.F.R. Ellis and M.A.H. MacCallum: *Comm. Math. Phys.* **12** (1969) 108.
- [43] M.P.Jr. Ryan and L.C. Shepley: *Homogeneous Relativistic Cosmology*, Princeton University Press, Princeton (1975).
- [44] U. Camci, I. Yavuz, H. Baysal, I. Tarhan and I. Yilmaz: *Ast. Sp. Sc.* **275** (2001) 391.
- [45] A. Einstein and W.de Sitter: *Proceeding of the National Academy of Sciences of the U.S.A.* **18** (1932) 213.
- [46] S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure of Space-time*, Cambridge University Press, Cambridge (1973).